arXiv:1212.3275v2 [quant-ph] 14 Dec 2012

Purified discord and multipartite entanglement

Eric G. Brown,[1] Eric J. Webster,[2] Eduardo Martín-Martínez,[2,3,4] and Achim Kempf[2,3,4]
[1]*Department of Physics and Astronomy, University of Waterloo, Waterloo, Ontario N2L 3G1, Canada*
[2]*Department of Applied Mathematics, University of Waterloo, Waterloo, Ontario, N2L 3G1, Canada*
[3]*Institute for Quantum Computing, University of Waterloo, Waterloo, Ontario, N2L 3G1, Canada*
[4]*Perimeter Institute for Theoretical Physics. Waterloo, Ontario N2L 2Y5, Canada*

We study the relationship between bipartite quantum discord in absence of entanglement and tripartite entanglement in the purified system. For qubits, we find that quantum discord requires the presence of both bipartite and genuinely tripartite quantum entanglement in the purified system. We consider generalizations of this result which then also allow us to explore the link between entanglement, discord and classical correlations in Gaussian states. Finally, we devise a protocol to remotely control the quantum discord resources available in a bipartite system AB.

I. INTRODUCTION

Quantum computation is believed to outperform its classical counterpart in its information processing efficiency. The resource most commonly associated with this improved performance is quantum entanglement. Operationally, entangled states are those which cannot be prepared through local operations and classical communication [1]. Equivalently, they are those states which cannot be written in the form $\sum_k p_k \rho_A^k \otimes \rho_B^k$, i.e. not separable. Entanglement is, however, difficult to create and manipulate in practice and this therefore greatly limits its theoretical benefits.

Recently it has become understood that entanglement is not the only correlation-resource capable of allowing a quantum speed-up. There exist quantum computational models which involve mixed, separable states such as the deterministic quantum computation with one qubit (DCQ1) [2]. The resource believed to provide an enhancement in these contexts is the so-called *quantum discord*, denoted $D(A,B)$. Discord is thought of as a generalized measure of quantum correlations, and therefore includes entanglement as a subset [3]. The nullity condition for discord $D(A,B)$ is more restricted than that for entanglement; specifically $D(A,B)=0$ iff the state can be written in the form $\sum_k p_k \rho_A^k \otimes |k\rangle_B \langle k|$, where $\{|k\rangle_B\}$ form an orthogonal set. A similar condition holds for the nullity of $D(B,A)$, and in general one can be zero while the other finite. It is known that for pure states, the discord becomes a measure of entanglement, more specifically, the entanglement entropy. For mixed states, however, it is not completely clear how discord is related to entanglement.

Given that discord without entanglement can only arise in a mixed state, we will here examine discord in terms of the entanglement found in its purification. We will find that, at least for qubit systems, 2-partite discord is intimately related to tri-partite entanglement in the purified system. If this finding generalizes, it may open the door to perhaps understanding n-partite entanglement in terms of (n-1) partite entanglement plus discord. This in turn could provide valuable new measures of discord and multi-partite entanglement and relationships between them. For example, we here find a close connection between the asymmetry of bi-partite discord and entanglement monogamy. We also propose a protocol to remotely increase quantum correlations of purely discord type between two distant parties.

Concretely, let us consider the simple scenario of pure, 3-qubit states $|\psi\rangle_{ABC}$ such that the reduced AB system, ρ_{AB}, is separable. Our goal is to understand the presence of quantum discord in ρ_{AB} in terms of the entanglement structure of $|\psi\rangle_{ABC}$. There are several motivations behind this. Firstly, since in the case of pure states the discord is equivalent to the entanglement entropy we should expect that the presence of discord in ρ_{AB}, despite the lack of entanglement, must arise via entanglement in its purification $|\psi\rangle_{ABC}$. Secondly, as we already mentioned, if we can indeed find a link between discord and purified entanglement then this may lead to a better understanding of and/or new measures of multipartite entanglement. The reason we choose to restrict ρ_{AB} to be separable is that, were it entangled, it would also be trivially discordant and we would not be able to attribute the presence of discord to entanglement in its purification. By restricting to separable states we are able to identify necessary and sufficient conditions on the purified entanglement in order for ρ_{AB} to have nonzero discord. Interestingly, we find that the presence of discord requires the purification to contain both bipartite *and* genuine tripartite entanglement.

In Sec. II we give a proof of these conditions for the previously discussed 3-qubits $|\psi\rangle_{ABC}$. Although limiting to 2-dimensional systems may seem quite restricted, we are also able to show that one of the conditions holds in general. In Sec. III we use these findings to motivate the investigation of discord and purified entanglement in the class of Gaussian continuous-variable states. In particular, we will use our findings to shed light on a quite remarkable property of discord in Gaussian states: that it is zero if and only if we have a product state – which means that there are no correlations (classical or quantum) present. We will confirm that the results found in Sec. II also hold for Gaussian states, or at least for the restricted class that we consider.

Finally, inspired by our findings in the former sections, in Sec. IV we devise a protocol to allow remote control over quantum resources, in the form of discord, shared by two partners. Namely, we consider two qubits A and B with no quantum correlations whatsoever between them. One is given to Alice (A) and the other is sent to Bob (B). If Alice does not want Bob to make use of quantum resources before she decides to authorize communication, it would be very convenient for Alice to have a means to 'remotely activate' quantum correlations between her and Bob's qubit, but without access to the latter. To achieve this, Alice is given another purifying qubit C that shares tripartite entanglement with A and B. We find that by acting on the AC system Alice is able to introduce quantum discord between A and B, despite there being no initial quantum correlation and with Alice having no access to B. In Sect. V we provide some concluding remarks.

II. ENTANGLEMENT STRUCTURE AND DISCORD

In this section we present our primary result, which will then be used to motivate the discussions in later sections. Here we restrict ourselves to the very simple system of a pair of qubits that are in a rank-2 state ρ_{AB}, such that its purification $|\psi\rangle_{ABC}$ will consist of only one extra qubit C. We further require the state ρ_{AB} to be separable. This is because we want to understand how the presence of discord between A and B is to be understood in terms of the entanglement structure of the purified system. If we were to allow entanglement between between A and B then this would trivially imply nonzero discord and there would therefore be no necessary conditions on the purified system to ensure the presence of discord.

To this end we can write the most general (up to relative phases) two-qubit, rank-2, separable state as

$$\rho_{AB} = q\,|0\rangle\langle 0|\otimes|0\rangle\langle 0| + (1-q)\,|\alpha\rangle\langle\alpha|\otimes|\beta\rangle\langle\beta|\,, \quad (1)$$

where $0 < q < 1$ and

$$|\alpha\rangle \equiv \cos\alpha\,|0\rangle + \sin\alpha\,|1\rangle\,, \quad (2)$$

$$|\beta\rangle \equiv \cos\beta\,|0\rangle + \sin\beta\,|1\rangle \quad (3)$$

are real combinations of the basis states $|0\rangle$ and $|1\rangle$. We don't consider the cases when $q = \{0, 1\}$ because then ρ_{AB} will be a (pure) product state and thus will trivially have zero discord. Note that we don't lose any generality by choosing the first projector to be $|0\rangle\langle 0|\otimes|0\rangle\langle 0|$. Also note however that we *have* lost generality by not including a relative phase between the two terms and by assuming that $|\alpha\rangle$ and $|\beta\rangle$ are *real* combinations of the basis vectors. This exclusion will not affect the primary result presented here, as will be explained below.

We now ask under what circumstances ρ_{AB} contains discord. Recall that the discord $D(A,B)$ when B performs the required measurement is not generally equivalent to the discord $D(B,A)$ when A performs the measurement. Indeed one can be zero while the other is nonzero. Clearly in our state of interest both of the discords will be trivially zero if $|\alpha\rangle = \pm|0\rangle$ or $|\beta\rangle = \pm|0\rangle$ because in this case ρ_{AB} is a product state. Aside from this we know that $D(A,B) = 0$ identically if $\langle\beta|0\rangle = 0$ [4], i.e. if $|\beta\rangle = \pm|1\rangle$. A similar condition holds for $D(B,A) = 0$. Concisely we can state

$$D(A,B) = 0$$
$$\text{iff } \{|\beta\rangle = \pm|1\rangle \quad \text{or} \quad |\alpha\rangle = \pm|0\rangle \quad \text{or} \quad |\beta\rangle = \pm|0\rangle\}, \quad (4)$$

and

$$D(B,A) = 0$$
$$\text{iff } \{|\alpha\rangle = \pm|1\rangle \quad \text{or} \quad |\alpha\rangle = \pm|0\rangle \quad \text{or} \quad |\beta\rangle = \pm|0\rangle\}. \quad (5)$$

The goal is now to compare these possibilities with those of the entanglement structure of the purification $|\psi\rangle_{ABC}$. Without loss of generality the purification is

$$|\psi\rangle_{ABC} = \sqrt{q}\,|0\rangle|0\rangle|0\rangle + \sqrt{1-q}\,|\alpha\rangle|\beta\rangle|1\rangle \quad (6)$$

Let us now consider from this the reduced state ρ_{AC}. Tracing over B and labeling $c_\alpha \equiv \cos\alpha$ etc, we have

$$\rho_{AC} =$$
$$\begin{pmatrix} q & \sqrt{q(1-q)}c_\beta c_\alpha & 0 & \sqrt{q(1-q)}c_\beta s_\alpha \\ \sqrt{q(1-q)}c_\beta c_\alpha & (1-q)c_\alpha^2 & 0 & (1-q)c_\alpha s_\alpha \\ 0 & 0 & 0 & 0 \\ \sqrt{q(1-q)}c_\beta s_\alpha & (1-q)c_\alpha s_\alpha & 0 & (1-q)s_\alpha^2 \end{pmatrix}. \quad (7)$$

The partially transposed eigenvalues of ρ_{AC} can then be readily computed and it is found that only one of the four, which we will call λ, can ever be negative. Recall that since ρ_{AC} has dimension 2×2 the positive partial transpose criterion is both a necessary and sufficient condition for separability [5]. Thus ρ_{AC} is entangled iff λ is negative. From this it is easy to show that ρ_{AC} is almost always entangled, being separable iff $\alpha = \{0, \pi\}$ or $\beta = \{\pi/2, 3\pi/2\}$, i.e. iff $|\alpha\rangle = \pm|0\rangle$ or $|\beta\rangle = \pm|1\rangle$. Additionally it becomes trivially separable if q is equal to 0 or 1, but we will not consider this case. The separability of ρ_{BC} follows similar conditions. Concisely:

$$\rho_{AC} \;\text{ is separable iff }\; \{|\alpha\rangle = \pm|0\rangle \quad \text{or} \quad |\beta\rangle = \pm|1\rangle\}, \quad (8)$$

$$\rho_{BC} \;\text{ is separable iff }\; \{|\beta\rangle = \pm|0\rangle \quad \text{or} \quad |\alpha\rangle = \pm|1\rangle\}. \quad (9)$$

Let us further consider the separability of ρ_{AC}, because the two different conditions for separability mean two very different things. We see that the first condition, $|\alpha\rangle = \pm|0\rangle$, coincides with the A system being a product onto the BC system; under this condition the 3-qubit state takes the product form $|\psi\rangle_{ABC} = |0\rangle\otimes|\psi\rangle_{BC}$. Of course in this case ρ_{AB} is separable, as we've seen, but it also clearly has zero discord and we see that this is compatible with Eqs. (4,5). The second condition, $|\beta\rangle = \pm|1\rangle$, is much more interesting. In this case the reduced state of A is mixed rather than pure, meaning that

$\|\alpha\rangle$	$\pm\|0\rangle$	$\neq\pm\|0\rangle$	$\pm\|0\rangle$	$\pm\|1\rangle$	$\neq\pm\{\|0\rangle,\|1\rangle\}$	$\neq\pm\{\|0\rangle,\|1\rangle\}$	$\pm\|1\rangle$
$\|\beta\rangle$	$\pm\|0\rangle$	$\pm\|0\rangle$	$\neq\pm\|0\rangle$	$\neq\pm\{\|0\rangle,\|1\rangle\}$	$\pm\|1\rangle$	$\neq\pm\{\|0\rangle,\|1\rangle\}$	$\pm\|1\rangle$
Entanglement structure							
$D(A,B)$	0	0	0	>0	0	>0	0
$D(B,A)$	0	0	0	0	>0	>0	0

FIG. 1. The relationship between the entanglement structure of $|\psi\rangle_{ABC}$ and the discord in ρ_{AB}. For given conditions on $|\alpha\rangle$ and $|\beta\rangle$ we display the resulting entanglement structure and the results for the discords $D(A,B)$ and $D(B,A)$. In the structure diagrams an ellipse represents the presence of bipartite entanglement while a triangle represents the presence of tripartite entanglement.



FIG. 2. The behavior of discord and bipartite negativity as we move in the (α,β) plane along a trajectory of constant tripartite entanglement $\pi_{ABC}=0.2$. $\varphi\in[0,2\pi)$ is a variable used to parameterize the trajectory through (α,β) space.

despite A not being entangled with B nor with C it *is* entangled with the BC system as a whole. In fact, in this case we have genuine tripartite entanglement occurring in the state $|\psi\rangle_{ABC}$, as can be confirmed by computing its π-tangle [6].

We can take the conditions for zero discord and separability in Eqs. (4,5) and (8,9) respectively and find that they match up very nicely; we display the resulting pattern in Fig. 1. There are two interesting things to notice from this pattern. Firstly, at least in the simple setting we are considering here, it appears that the presence of discord in ρ_{AB} requires the presence of *both* bipartite entanglement *and* tripartite entanglement in the purified system (notice that when there is no bipartite entanglement, i.e. a GHZ type state $|000\rangle+|111\rangle$, there is no discord). Secondly we very clearly see the asymmetry of discord $D(A,B)\neq D(B,A)$ represented in the entanglement structure.

As an important note, recall that we neglected to include relative phases in our state (1). In the analysis we've done here this is not a problem and the structure in Fig. 1 will continue to hold if phases are included. The reasons for this are that 1) the nullity of discord depends only on the orthogonality the projectors in Eq. (1), which for us will not be affected by phases and 2) the partially transposed eigenvalues do not depend on any relative phases and thus the entanglement structure of $|\psi\rangle_{ABC}$ will be independent of them as well. Thus we are justified in using the simplifying assumption of no phases, reducing our phase space from six dimensions (q, α and β plus three phases) down to three dimensions.

As an aside, it is interesting to examine how the discords $D(A,B)$, $D(B,A)$ quantitatively relate to the negativities $\mathcal{N}_{AC}$, $\mathcal{N}_{BC}$ and the π-tangle π_{ABC}. The structure seen in Fig. 1 would seem to suggest that an increase in $D(A,B)$ (that is, considering two different states of the form (6) with differing discord) should be accompanied either by an increase in $\mathcal{N}_{AC}$ or in π_{ABC}, or both. Similarly $D(B,A)$ seems that it should increase with increasing $\mathcal{N}_{BC}$ or π_{ABC}, or both. To test this we set $q=1/2$ and we consider two trajectories through the (α,β) plane characterizing our state. One of these is a path of constant π_{ABC} and the other is a path of constant $\mathcal{N}_{AC}$. We then plot the remaining quantities as functions of an arbitrary parameter φ that parameterizes these paths. This will let us, for example, deduce how $D(A,B)$ changes as $\mathcal{N}_{AC}$ changes but while π_{ABC} is kept at a constant, nonzero value.

These plots are displayed in Fig. 2, where we keep constant $\pi_{ABC}=0.2$, and in Fig. 3, where we keep constant $\mathcal{N}_{AC}=0.1$. In Fig. 2 we observe the behavior that was expected, namely $D(A,B)$ is perfectly monotonic with $\mathcal{N}_{AC}$ and $D(B,A)$ is perfectly monotonic with $\mathcal{N}_{BC}$. In Fig. 3 however we find something rather different, namely we find that that neither $D(A,B)$ nor $D(B,A)$ is always monotonic with π_{ABC}. This is not surprising in the case of $D(B,A)$ because, as we see, the negativity $\mathcal{N}_{BC}$ (which we know feeds $D(B,A)$) experiences a dra-



FIG. 3. The behavior of discord, negativity, and π-tangle as we move in the (α, β) plane along a trajectory of constant bipartite entanglement $\mathcal{N}_{AC} = 0.1$. $\varphi \in [0, 2\pi)$ is a variable used to parameterize the trajectory through (α, β) space.

matic decrease. It is surprising, however, that we also see a decrease in $D(A, B)$ during this period, despite $\mathcal{N}_{AC}$ remaining constant and π_{ABC} increasing. Evidently, while $\mathcal{N}_{BC}$ does not play a role in the nullity of $D(A, B)$, as seen from Fig. 1, it does generally contribute to its value.

There is another interesting observation that can be made from Fig. 2: For a fixed value of purely tripartite correlations, since discord $D(A, B)$ increases with entanglement $\mathcal{N}_{AC}$ and $D(B, A)$ increases with entanglement $\mathcal{N}_{BC}$, we notice that the fundamental asymmetry between $D(A, B)$ and $D(B, A)$ stems from the entanglement monogamy principle ($\mathcal{N}_{AC}$ and $\mathcal{N}_{BC}$ are anticorrelated).

Although we have shown the validity of the structure seen in Fig. 1 in only a very restricted system and class of states, namely rank-2 states of a two-qubit system, it is tempting to postulate that such a structure holds for more general systems as well. In fact, it is easy to show that two of the statements hold true in general: if AC is separable, then $D(A, B) = 0$ and if BC is separable, then $D(B, A) = 0$. To show this, recall the expression for $D(A, B)$ [7]

$$D(A, B) = E_{AC} + S_B - S_C,$$

where E is the entanglement of formation and S the von Neumann entropy. The corresponding expression for $D(A, C)$ is

$$D(A, C) = E_{AB} + S_C - S_B.$$

Together, these yield

$$D(A, B) + D(A, C) = E_{AC} + E_{AB},$$

but in our case $E_{AB} = 0$ to begin with, and if AC is also separable then $D(A, B) + D(A, C) = 0 \rightarrow D(A, B) = 0$, since discord is always non-negative. A similar argument can be made for the BC separable case.

III. UNDERSTANDING THE VANISHING OF DISCORD IN GAUSSIAN STATES

So far, we have shown the structure in Fig. 1 for 3-qubit systems. The validity of two of the requirements for such a structure in general motivates us to look for other classes of states for which this structure holds. To this end we consider a restricted class of Gaussian states and find that the structure in Fig. 1 holds true. This finding also sheds light on a peculiar property of quantum discord in Gaussian states. In particular we consider the quantum discord in two-mode Gaussian states, which are known to have the property that the discord vanishes if and only if the state is a product state, as first suggested in [8] and later proven in [9]. This is a rather surprising result, especially in light of the fact that Gaussian states are often considered to be the most classical of quantum states. It seems however that in such states it is impossible to banish quantum correlations without also eliminating any trace of classical correlations. We will find that, again for a rather restricted class of Gaussian states, this property is consistent with the entanglement structure of the purified system. This finding not only provides support for the generality of the structure in Fig. 1, but also gives insight into this strange property of Gaussian states.

We will first give a very brief introduction to Gaussian states, focusing only on the concepts necessary for our purposes here. For a more complete introduction the reader is referred to [10, 11], among many other resources available in the literature. In short, such states are an important class of continuous-variable states for which the characteristic phase space functions are Gaussian. This means that they can be fully characterized by their first and second quadrature moments, which are typically arranged in the *covariance matrix*, the elements of which for an N-mode state are $\sigma_{(ij)} \equiv \langle \hat{X}_i \hat{X}_j + \hat{X}_j \hat{X}_i \rangle - 2\langle X_i \rangle \langle X_j \rangle$ where $\hat{\boldsymbol{X}} = (\hat{x}_1, \hat{p}_2, \ldots \hat{x}_N, \hat{p}_N)$ is a the vector containing the $2N$ quadrature operators. For a two-mode Gaussian state ρ_{AB} the covariance matrix is of the form

$$\sigma_{AB} = \begin{pmatrix} \alpha & \gamma \\ \gamma^T & \beta \end{pmatrix}, \tag{10}$$

where α and β are the 2×2 covariance matrices corresponding to the reduced states $\rho_A = \mathrm{Tr}_B \rho_{AB}$ and $\rho_B = \mathrm{Tr}_A \rho_{AB}$, which are guaranteed to also be Gaussian. The matrix γ encodes the correlations between A and B, both quantum and classical. Higher-mode states are similarly defined. Note that finding reduced states is trivially easy in this formalism; one simply neglects the rows and columns associated with the modes one chooses to trace over. For two mode Gaussian states, like two-qubit states, we have a necessary and sufficient PPT criterion for determining separability that reads [12]

$$\Delta_{AB} \leq 1 + \det\sigma_{AB}, \tag{11}$$

where $\Delta_{AB} = \det\alpha + \det\beta - 2\det\gamma$.

Unitary transformations in the Hilbert space $\rho \to U\rho U^\dagger$ are represented in the phase space picture by *symplectic transformations* $\sigma \to S\sigma S^T$, where the matrix S is required to satisfy $S\Sigma S^T = S^T\Sigma S = \Sigma$ and where for N-modes Σ is the symplectic form

$$\Sigma = \bigoplus_{i=1}^{N} \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix}. \tag{12}$$

For any N-mode Gaussian state we can find a symplectic transformation that brings the covariance matrix into the diagonal form [13] (the so called Williamson normal form) $\sigma = \mathrm{diag}(\nu_1, \nu_1, \nu_2, \nu_2, \ldots \nu_N, \nu_N)$, where the numbers ν_i are the state's *symplectic eigenvalues* which can be computed as the eigenvalues of the matrix $|i\Sigma\sigma|$.

In Gaussian states the uncertainty principle takes the form $\nu_i \geq 1$ for all i. Furthermore the symplectic eigenvalues encode the mixedness of the state, where $\nu_i = 1$ for all i corresponds to a pure state, and more generally the purity of a state is given by $\mathrm{Tr}\rho^2 = \prod_i \nu_i^{-1}$. Note therefore that pure Gaussian states saturate the uncertainty relation, and more generally the uncertainty in a state above that imposed by the uncertainty principle is due only to possible mixedness, corresponding to classical uncertainty. This is why Gaussian states are often considered the "most classical" of quantum states.

Our goal is to consider two mode Gaussian states and their purification in the hopes that we can find consistency, similar to Fig. 1, between the fact that quantum discord in Gaussian states vanishes only for product states and the purified entanglement structure. Through the action of local symplectic transformations we can always cast a two-mode covariance matrix into the form

$$\sigma_{AB} = \begin{pmatrix} a & 0 & c & 0 \\ 0 & a & 0 & d \\ c & 0 & b & 0 \\ 0 & d & 0 & b \end{pmatrix}, \tag{13}$$

where a and b are the symplectic eigenvalues of the reduced states ρ_A and ρ_B, respectively. This state is a product state if and only if $c = d = 0$.

Similar to qubits, we will constrain ourselves to a restricted class of states. Namely, we will consider separable states that can be purified with only a single extra mode. Regarding the separability condition, for computational simplicity we will further restrict ourselves to states for which $c = d$. This condition guarantees separability, but it does not account for all separable states. Given this, our necessary and sufficient criterion for σ_{AB} being a product state is $c = 0$. Note that if $a = 1$ or $b = 1$ we also have a product state, since this corresponds to ρ_A or ρ_B being pure, however in this case the uncertainty relation reduces to $c = 0$, so these conditions for having a product state are in fact equivalent. In order for σ_{AB} to be purified with one additional mode we also require that one of its two symplectic eigenvalues be equal to unity; let this be $\nu_1 = 1$. This is equivalent to the condition of the qubit state being rank-2 from the previous section. Imposing these restrictions and computing the symplectic eigenvalues we are able to find the equalities

$$b = \frac{c^2 + a - 1}{a - 1}, \quad \nu_2 \equiv \nu = \frac{c^2 + a(a-1)}{a-1}. \tag{14}$$

That is, the condition $\nu_1 = 1$ along with $c = d$ reduces our system to only two free parameters, which we have chosen to be a and c.

In order to perform the purification we must first put the two-mode state into its Williamson normal form, namely $\sigma_{AB}^{(W)} = \mathrm{diag}(1, 1, \nu, \nu)$. This can be performed by a symplectic transformation on the AB system. In particular, we find the symplectic matrix that satisfies $S_{AB}\sigma_{AB}^{(W)}S_{AB}^T = \sigma_{AB}$ is of the form

$$S_{AB} = \frac{1}{\sqrt{c^2 + (a-1)^2}} \begin{pmatrix} -c & 0 & a-1 & 0 \\ 0 & -c & 0 & a-1 \\ a-1 & 0 & c & 0 \\ 0 & a-1 & 0 & c \end{pmatrix}. \tag{15}$$

The purified state $|\psi\rangle_{ABC}$, involving three modes ABC, has a covariance matrix that is given in a form essentially equivalent to a Schmidt decomposition [14]:

$$\sigma_{ABC}^{(S)} = \begin{pmatrix} \sigma_{AB}^{(W)} & \gamma_\nu \\ \gamma_\nu^T & \sigma_C^{(W)} \end{pmatrix}, \tag{16}$$

where $\sigma_C^{(W)} = \mathrm{diag}(\nu, \nu)$ is the covariance matrix of the reduced C state and

$$\gamma_\nu = \begin{pmatrix} 0 & 0 \\ 0 & 0 \\ \sqrt{\nu^2 - 1} & 0 \\ 0 & -\sqrt{\nu^2 - 1} \end{pmatrix}. \tag{17}$$

Note that, as must be the case, the mixedness of the mode C is equivalent to that of the two-mode AB state. If in doubt, one can easily compute the symplectic eigenvalues of $\sigma_{ABC}^{(W)}$ and confirm that they are all unity, meaning that the three-mode state is pure. Note also that if $\nu = 1$, meaning that ρ_{AB} and ρ_C are pure, then the correlation matrix γ_ν vanishes as it must.

In order to examine the entanglement structure in the purified system, namely the entanglement in the states ρ_{AC} and ρ_{BC}, we must reverse the symplectic transformation that we used to put σ_{AB} into normal form since in doing so we have mixed the modes A and B. That is, we need the ABC state in the form

$$\begin{aligned}\sigma_{ABC} &= (S_{AB} \oplus I_C)\sigma^{(S)}_{ABC}(S^T_{AB} \oplus I_C) \\ &= \begin{pmatrix} S_{AB}\sigma^{(W)}_{AB}S^T_{AB} & S_{AB}\gamma_\nu \\ (S_{AB}\gamma_\nu)^T & \sigma^{(W)}_C \end{pmatrix} \\ &= \begin{pmatrix} \sigma_{AB} & S_{AB}\gamma_\nu \\ (S_{AB}\gamma_\nu)^T & \sigma^{(W)}_C \end{pmatrix}.\end{aligned} \tag{18}$$

With this matrix in hand we are easily able to compute the covariance matrices σ_{AC} and σ_{BC} by simply eliminating the rows and columns associated with B and A respectively. We can then check the separability of these states using the criterion in Eq. (11). For both reduced states it is easy to show that they are separable if and only if $c = 0$, in which case Eq. (11) is saturated. That is, ρ_{AC} and ρ_{BC} are separable if and only if ρ_{AB} is a product state. We can similarly check the presence of genuine tripartite entanglement in the ABC system. For this we will use the measure of tripartite entanglement known as the *residual Gaussian cotangle* [15]. We will not include its rather unsightly form here, however the reader may find it on Pg. 8 of [15]. The result is that the tripartite entanglement between A, B and C similarly vanishes if and only if $c = 0$.

In summary, the bipartite entanglement in the AC and BC systems, as well as the tripartite entanglement in the pure ABC system, each vanish if and only if the AB system is a product state. We also know that in Gaussian states the quantum discord between A and B vanishes if and only if it is a product state. We have thus found that Gaussian states, or at least the restricted set of Gaussian states that we have considered, satisfy the relation between discord and purified entanglement displayed in Fig. 1. This provides some partial support for the possibility of this structure being a general property of quantum states, and furthermore it provides an alternative route to understanding the surprising result that discord in Gaussian states vanishes only for product states.

IV. PROTOCOL FOR REMOTE ACTIVATION OF DISCORD

Taking the result found in Section II as inspiration, here we devise a simple protocol to locally increase discord between two distant parties that initially have no quantum correlations. This would be useful in a scenario in which Alice needs to share quantum resources with a distant partner Bob but she does not want Bob to be able to utilize those resources unless previously authorized by Alice.

It is known that it is possible to remotely activate entanglement in this sense: Assume that Alice and Bob are seperated by a large distance and initially do not share any quantum correlations. Further assume that Bob is entangled with a system C. Then, Alice may ‘activate’ entanglement between her and Bob by locally interacting with C. For example, Alice and Bob can be chosen to be Unruh deWitt detectors, system C can be chosen to be the vacuum state, and a suitable interaction protocol is then that of entanglement swapping from the vacuum [16–21]. Notice that no tri-partite entanglement is required here. It is not even necessary for A, C and B, C to be entangled. Namely, as was shown in [22], A, B entanglement can be established withouit prior entanglement, namely if A, C and B, C are merely discorded - and if in addition the crucial condition holds that A, B started out discorded.

Now, let us ask under which conditions, if Alice and Bob start without *any* quantum correlations, Alice may ‘activate’ A, B discord with Bob. Interestingly, we find that this is possible and it requires only tri-partite A, B, C entanglement.

To see this, let us consider two partners Alice (who possesses some qubit A and an ancillary qubit C) and Bob (who has a qubit B), such that A and B have no quantum correlations. Recalling Fig. 2, we see that if the ABC system is prepared to have tripartite entanglement then if Alice is able to introduce entanglement between A and C without destroying the tripartite entanglement this will produce discord between A and B. Since Alice has access to both A and C she is indeed able to inject such correlations into the system via a unitary operation. The advantage of this protocol is that Alice can remotely deny or allow any quantum information tasks that may be performed between her and Bob making use of these resources.

To demonstrate this explicitly we start by preparing a three qubit state of the form

$$|\psi\rangle_{ABC} = \sqrt{q}\,|000\rangle + \sqrt{1-q}\,|111\rangle\,, \tag{19}$$

which is separable between A and B. It also has zero quantum discord $D(A, B)$ which we plan to increase. As usual we omit $q = 0, 1$ as these cases are trivial. We also omit $q = 1/2$ because entanglement AC is zero when ABC is maximally entangled. In addition to AB being separable, all other bipartite subsystems of this state are also separable and the π-tangle is $\pi_{ABC} = q(1-q)$. Assuming we can keep π_{ABC} constant then we expect discord $D(A, B)$ to be monotonically correlated with entanglement AC, as indicated by Fig. 2. As such, maximizing the increase in entanglement between A and C will maximize the resulting discord between A and B.

For example, consider acting on $|\psi\rangle_{ABC}$ with the unitary $U_{AC} \otimes I_B$, where

$$U_{AC}(t) = \begin{pmatrix} \cos t & 0 & 0 & -\sin t \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ \sin t & 0 & 0 & \cos t. \end{pmatrix}$$

This transformation leaves entanglement AB, BC and ABC unchanged, while modifying entanglement AC. Note however that this statement is only correct when our state is of the form Eq. (19); more generally entanglement in AB, BC and ABC will also be modified. This



FIG. 4. Discord $D(A,B)$ and entanglement AC following the unitary $U_{AC}(t)$ being applied to the AC system

said, the main result of this section, namely the ability to remotely activate quantum correlations, still holds in general. Plotting $\mathcal{N}_{AC}$ and $D(A,B)$ in Fig. 4 for the transformed state as a function of t confirms that the discord AB is totally correlated with the entanglement AC.

The Hamiltionian that generates this unitary evolution $U_{AC}(t) = e^{-i\mathcal{H}_{AC}t/\hbar}$ is given by

$$\mathcal{H}_{AC} = \hbar \begin{pmatrix} 0 & 0 & 0 & -i \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ i & 0 & 0 & 0, \end{pmatrix}$$

and in fact any Hamiltonian evolution on the AC system of our state $|\psi\rangle_{ABC}$ of the form

$$\mathcal{H}_{AC}^{X} = \hbar \begin{pmatrix} h_{00} & 0 & 0 & h_{03} \\ 0 & h_{11} & h_{12} & 0 \\ 0 & h_{12}^{*} & h_{22} & 0 \\ h_{03}^{*} & 0 & 0 & h_{33} \end{pmatrix}$$

will have the effect of leaving the π-tangle and all bipartite entanglements other than AC constant, and thus can be used to increase the discord $D(A,B)$ of $|\psi\rangle_{ABC}$. By controlling the degrees of freedom in $\mathcal{H}_{AC}^{X}$ we are able to modify the shape of the curves in Fig. 4 while maintaining monotonic correlation between $D(A,B)$ and $\mathcal{N}_{AC}$. By appropriately choosing the time and Hamiltonian of evolution mediating in the AC system Alice can thus have good control over the amount of discord that she injects between herself and Bob.

We have shown that we have a protocol for increasing discord between two parties, which initally have no discord, without acting at all on one of the parties. Tripartite entanglement is a necessary resource for this to work, but not sufficient because a maximally entangled $|\psi\rangle_{ABC}$ would not work. Not only is this an interesting and possibly useful protocol to increase discord, it also provides a link between discord and bipartite entanglement with the environment.

V. CONCLUSIONS

We have studied the relationship between the discord in an unentangled bipartite system A, B and the bipartite and tripartite entanglement found in its purification A, B, C.

We have found that both purely tripartite entanglement and bipartite entanglement between A, C (or B, C) are necessary to have non-zero discord $D(A,B)$ (or $D(B,A)$). We found that there is a trade-off between the two directions of discord between A and B. Both cannot be large at the same time because their strength relies on the strength of the A, C or B, C entanglement respectively. However, the A, C and B, C entanglements cannot be strong simultaneously because of entanglement monogamy. We have therefore shown that the asymmetry between $D(A,B)$ or $D(B,A)$ stems from entanglement monogamy.

This hierarchy of quantum correlations suggests that discord may be characterizable by means of different kinds of multipartite entanglement in an extended space where we consider the state AB and its environment. We thoroughly analyzed the case of two-dimensional systems and showed that some results, e.g., the relation between the asymmetry of discord and entanglement monogamy hold more generally.

We have also used our findings to obtain insight on the behaviour of correlations in Gaussian states reported in [8] and [9]. We have observed that, at least for a limited class of Gaussian states, the correlation hierarchy found in Sect. II does apply to these continuous-variable states.

Lastly, we have used the knowledge gained in previous sections to suggest a protocol to remotely control the discord quantum resources between two partners A and B through increasing bipartite correlation between A and a common environment or ancillary system in the background of tripartite entanglement.

Finally, an alternate way to interpret our results suggests further analysis: Assume that a system B is entangled with a composite system AC so that there is tripartite A, B, C entanglement and A, C entanglement but no A, B entanglement. Then, system AC breaks up in a way that leaves A and C entangled. Our results show that the following is possible: While B does not acquire entanglement with A from the breakup of the AC system, B can acquire discord with A in the breakup. In other words, tri-partite entanglement can manifest itself on the level of 2-partite quantum correlations as discord.

It is tempting to conjecture that discord will keep playing a central role in the relationship between n-partite and (n-1) partite entanglement.

To this end, it should be very interesting to generalize our strategy of purifying discord to more than three and also larger quantum systems. In fact, it should be possible and very interesting, also for practical purposes, to investigate the corresponding Hamiltonians, i.e., to study which types of interactions give rise to the structures of discord and its purification that we consider in this program.

VI. ACKNOWLEDGEMENTS

The authors gratefully acknowledge support through NSERCs Banting, Discovery and Canada Research Chairs Programs.

[1] R. F. Werner, Phys. Rev. A **40**, 4277 (1989).

[2] E. Knill & R. Laflamme, Phys. Rev. Lett. **81**, 5672-5675 (1998).

[3] H. Ollivier & W. H. Zurek, Phys. Rev. Lett. **88**, 017901 (2001).

[4] K. Modi *et. al.*, arXiv:1112.6238 [quant-ph]

[5] A. Peres, Phys. Rev. Lett. **77**, 1413 (1996); M. Horodecki, P. Horodecki and R. Horodecki, Phys. Lett. A. **223**, 1 (1996)

[6] Y. Ou and H. Fan, Phys. Rev. A. **75**, 062308 (2007)

[7] Felipe F. Fanchini, *et. al.*, arXiv:1006.2460v3 [quant-ph] (2011)

[8] G.Adesso and A. Datta, Phys. Rev. Lett. **105**, 030501 (2010).

[9] S. Rahimi-Keshari, C. M. Caves and T. C. Ralph, arXiv:1209.2221 [quant-ph]

[10] B. L. Schumaker, Physics Reports **135**, 317 (1986)

[11] G. Adesso and F. Illuminati, J. Phys. A: Math. Theor. **40**, 7821 (2007)

[12] A. Serafini, F. Illuminati and S. D. Siena, J. Phys. B: At. Mol. Opt. Phys. **37**, L21 (2004)

[13] J. Williamson, Am. J. Math. **58**, 141 (1936)

[14] A. Botero and B. Reznik, Phys. Rev. A. **67**, 052311 (2003)

[15] G. Adesso and F. Illuminati, New. J. Phys. **8**, 15 (2006)

[16] B. Reznik, A. Retzker, and J. Silman, Phys. Rev. A **71**, 042104 (2005)

[17] G. VerSteeg and N. C. Menicucci, Phys. Rev. D **79**, 044027 (2009).

[18] C. Sabin, J. J. Garcia-Ripoll, E. Solano, J. Leon Phys. Rev. B **81**, 184501 (2010)

[19] M. Cliche and A. Kempf, Phys. Rev. D **83**, 045019 (2011)

[20] S. Jay. Olson and Timothy C. Ralph, Phys. Rev. Lett. **106**, 110404 (2011)

[21] C. Sabin, B. Peropadre, M. Del Rey, E. Martín-Martínez Phys. Rev. Lett. **109**, 033602 (2012)

[22] T. K. Chuan, J. Maillard, K. Modi, T. Paterek, M. Paternostro,M. Piani, Phys. Rev. Lett. **109**, 070501 (2012)